

Abortion in Canada

Lessons learned and opportunities for expanding access to care

Angel M. Foster, DPhil, MD, AM
She/her/hers/anything respectful
Professor, Faculty of Health Sciences
2011-2016 Endowed Chair in Women's Health Research
University of Ottawa

Outline for today

- Brief history of abortion in Canada
- Abortion in Canada today
 - Legal status
 - Prevalence and service delivery
 - The role of medication abortion
- Key learnings from the journey of abortion in Canada



1869

Abortion is made illegal





1869

Abortion is made illegal



1969

Criminal code amended



1869

Abortion is made illegal

1970s-1980s

Defiance & mobilization



1969

Criminal code amended




Abortion is made illegal

1970s-1980s
Defiance & mobilization


1969
Criminal code R. v. Morgentaler amended







Mifepristone's journey in Canada





Abortion in Canada today




- Abortion does not live in the criminal code
 - No gestational age restrictions
 - No bans or requirements with respect to reason
- Abortion is considered a medically necessary service
 - Instrumentation and medication abortion are covered by provincial, territorial, and federal insurance schemes
 - Adolescents are able to obtain abortion care without parental involvement (with some exceptions)
 - No partner or spousal notification or consent is required
- Abortion is regulated at the provincial, institutional, and professional college/association levels



Location, location, location

No clinic services are available in the territories: All interritory abortions are performed in hospitals

Women in rural areas may have to go to five consultations to have an abortion at a facility



In 2017, on-Island services offered for the first time in 35 years

Clinic-based instrumentation abortions not covered by provincial insurance



Abortion by the numbers



- One in three Canadian women will have an abortion over the course of their reproductive lives
- □ 100,000+ abortions performed each year
 - Free-standing clinics, hospitals, primary care clinicians
- 90% of abortions in the first trimester
 - About 40% of all abortions are with mifepristone/misoprostol
- Less than 1% of abortions take place after 20 weeks from the first day of the last menstrual period (LMP)
 - Almost all of the 100 abortions each year after 24 weeks take place in the US



Other challenges

- Abortion surveillance
- Training the next generation of providers
- Belief-based denial of care (conscientious objection)
- Protests, protective zones, and legal challenges
- The United States



Some key learnings from the abortion journey in Canada

- Decriminalization provides a different abortion framework but is not a panacea
 - Policies, regulations, health system structures, and broader systems of oppression create barriers to access
- Abortion can be fully integrated into a health system and normalized as a medically necessary service
 - Inequities in the health system can then impact abortion care
- Medication abortion care be provided safely and effectively through a variety of service delivery systems
 - Non-evidence-based regulations limit the potential of medication abortion to address considerable access barriers
- Abortion care after 20 weeks is critical
 - Continued attention and advocacy is required to ensure equitable access

Questions & discussion

angel.foster@uottawa.ca

